



Derived Cats,
Arithmetic, &
Rationality
UCSC 4/23

Alicia Samarche

University
of Utah

slides:

You can follow along with these slides by downloading them from

alicia.lamarche.xyz/talks/04232021.pdf

↖ Contains links to
references & more!

Forthcoming work with :

Matthew Ballard

University of South Carolina

matthewrobertballard.com

arXiv:?????.?????

also contains joint work with:

Matthew Ballard

University of South Carolina

matthewrobertballard.com

Alexander Duncan

University of South Carolina

people.math.sc.edu/duncan

Patrick McFaddin

Fordham University

mcfaddin.github.io

[arXiv:2006.06876](https://arxiv.org/abs/2006.06876)

&

[arXiv:2009.10175](https://arxiv.org/abs/2009.10175)

Question:

Can $D^b(X)$ be used to answer rationality questions about X ?

if not, why not?

if so,
to what extent?

X will denote a smooth projective variety throughout.


outline

1. Motivation
2. Derived Categories
3. Arithmetic Toric Varieties
4. Results

Notation. For a smooth projective variety X , $D^b(X) := D^b(\text{Coh}(X))$
//
bounded derived category of coherent sheaves on X

We say that a variety X defined over a field k has a ***rational point*** (or ***k -rational point***) if there is a map $\text{Spec}(k) \rightarrow X$.

Definition. A variety X defined over a field k is called **rational** if it is *birational* to $\mathbb{P}^n$ for some n .




How can you determine if two varieties are birational?

Theorem. (*Weak Factorization*) Any birational map between two smooth complex varieties can be decomposed as a series of finitely-many blow-ups/blow-downs along smooth subvarieties.

Naive wishlist for a tool to detect rationality:

- Is relatively ‘nice’ for $\mathbb{P}^n$
- Behaves with respect to weak factorization



Theorem. ([Beilinson](#)) $D^b(\text{Coh}(\mathbb{P}^n))$ ‘decomposes’ in the following way:

$$D^b(\text{Coh}(\mathbb{P}^n)) = \langle \mathcal{O}, \mathcal{O}(1), \dots, \mathcal{O}(n) \rangle$$

Theorem. ([Bondal/Orlov](#)) Let $Y \subset X$ a smooth subvariety of codimension c .

$$\begin{array}{ccc} E & \xrightarrow{i} & Bl_Y(X) \\ p \downarrow & & \downarrow \pi \\ Y & \xrightarrow{\quad} & X \end{array}$$

Then, we have the following decomposition of $D^b(Bl_Y(X))$:

$$D^b(Bl_Y(X)) = \langle \pi^* D^b(X), i_* p^* D^b(Y), i_* p^* D^b(Y) \otimes \mathcal{O}_p(1), \dots, i_* p^* D^b(Y) \otimes \mathcal{O}_p(c-2) \rangle$$

ideq:

X rational $\implies D^b(X)$ is not ‘too’ complicated?

$D^b(X)$ is not ‘too’ complicated $\implies X$ rational?

How do we ‘measure’ how complicated $D^b(X)$ is?

Conjecture. (Orlov) A smooth projective variety with a full exceptional collection is rational.

Conjecture. (Lunts) Over a general field k , a full k -exceptional collection for $D^b(X)$ implies that X admits a locally-closed stratification into subvarieties that are each k -rational.

Conjecture. (*Orlov*) A smooth projective variety with a full exceptional collection is rational.

Conjecture. (*Lunts*) Over a general field k , a full k -exceptional collection for $D^b(X)$ implies that X admits a locally-closed stratification into subvarieties that are each k -rational.

Theorem. ([*Ballard, Duncan, L., McFaddin*](#)) A smooth projective toric variety X over a field k with $X(k) \neq \emptyset$ possessing a full k -exceptional collection is rational.

Question. For X defined over a field k , if $D^b(X)$ admits a particular type of ‘nice’ decomposition, does this mean that X is rational over k ?

Evidence:

For a Severi-Brauer curve X , $D^b(X)$ can be decomposed, but is only ‘nice’ when X is trivial.

dimension 1



- Genus 0: nontrivial Severi-Brauer curves never have k -exceptional collections. (Unless they are trivial- i.e. $\mathbb{P}_k^1$)
- Genus ≥ 1 curves do not admit nontrivial semiorthogonal decompositions.



([Okawa, 2011](#))

Question. For X defined over a field k , if $D^b(X)$ admits a particular type of ‘nice’ decomposition, does this mean that X is rational over k ?

Evidence:

dimension 2

- [Vial \(2016\)](#)

{ full k -exceptional collection & geometrically rational }

$\implies$ rational

- [Brown & Shipman \(2015\)](#)

{ full strong exceptional collection of line bundles
(over $\mathbb{C}$) & stability conditions }

$\implies$ rational

Question. Can $D^b(X)$ be used to answer rationality questions about X ?

Question. If X is defined over a field k and $D^b(X)$ admits a particular type of ‘nice’ decomposition, does this mean that X is ~~rational~~ over k ?



stably rational
retract rational
unirational
...

Question. For X defined over a field k , if $D^b(X)$ admits a particular type of ‘nice’ decomposition, does this mean that X has a *rational point* over k ?

The derived category can be used to detect the difference between ‘generalized Del Pezzo varieties’ and their twists by torsors.

Theorem. (*Ballard, L.*) If X is a generalized Del Pezzo variety, then X has a rational point if and only if $D^b(X)$ admits a full étale exceptional collection.

Question. For X defined over a field k , can $D^b(X)$ detect the existence of k -rational points on X ?

Addington, Antieau, Frei, Honigs (2019)

Gave an example of two derived equivalent varieties where one has a rational point and the other does not.

~~1. Motivation~~

② Derived Categories

3. Arithmetic Toric Varieties

4. Results

Precise definitions of what follows can be found in
Ballard, Duncan, McFaddin (2017)

Definition. Let A be a finite dimensional k -algebra of finite homological dimension. An object E of $D^b(X)$ is **A -exceptional** if:

- $\text{End}(E) \cong A$
- $\text{Hom}(E, E[n]) = 0$ for all $n \neq 0$

An object E is **exceptional** if it is A -exceptional for a division algebra A , and is **étale exceptional** if A is a finite separable field extension of k .

A totally ordered set $\{E_1, \dots, E_s\}$ of exceptional objects is a **full exceptional collection** if

- Each E_i is A_i -exceptional for $A_i = \text{End}(E_i)$
- $\text{Hom}(E_i, E_j[n]) = 0$ for all n , whenever $i > j$
- The collection of E_i generate the category


Observation. A (triangulated) category can be decomposed as the derived categories of (smooth) points if and only if it possesses a full étale exceptional collection.

Question. Which varieties admit full exceptional collections?

Theorem. (Ballard, Duncan, McFaddin, [2017](#))

A k -variety X admits a full exceptional collection if and only if $X_{k^{sep}}$ admits a full exceptional collection whose components are permuted by the action of $\text{Gal}(k^{sep}/k)$.

Theorem. (Kawamata, [2006/2013](#)) For a smooth toric variety X over an algebraically closed field k of characteristic zero, $D^b(X)$ admits a full exceptional collection.



In general, determining if $D^b(X)$ admits a full exceptional collection for an arbitrary variety X is difficult.

~~1. Motivation~~

~~2. Derived Categories~~


3. Arithmetic Toric Varieties

4. Results

Definition. A **torus** over a field k is an algebraic group T such that $T_{k^{sep}} \cong \mathbb{G}_m^n$. We say that T is **split** if $T \cong \mathbb{G}_m^n$.

Definition. A **toric variety** is an algebraic variety that contains an algebraic torus as a dense open subset in such a way that the action of the torus on itself extends to the entire variety.




The geometry of a normal toric variety over a separably closed field is completely determined by the combinatorial data of its associated fan- a diagram that describes how the variety is constructed via the gluing of torus orbits.



Definition. An *arithmetic toric variety* over k is a variety that base-changes to a toric variety over k^{sep} .

“Twisted Forms of Toric Varieties” ([Duncan. 2014](#))

“Arithmetic Toric Varieties” ([Elizondo, Lima-Filho, Sottile, Teitler. 2010](#))



Example

$\mathbb{P}_{\mathbb{R}}^1$

$\mathbb{R}^\times$


acts via $\lambda \cdot [x : y] = [\lambda x : y]$

S^1


*acts via rotation matrices
(or complex multiplication)*

Example

$\mathbb{P}_{\mathbb{R}}^1$



Example:




Theorem. (*Ballard, L.*) If X is a generalized Del Pezzo variety, then X has a rational point if and only if $D^b(X)$ admits a full étale exceptional collection.

an arithmetic
toric variety!

The fan for V_{2n} has the following rays:

$$\begin{array}{ll} e_0 = (-1, \dots, -1) & \bar{e}_0 = (1, \dots, 1) \\ e_1 = (1, 0, \dots, 0) & \bar{e}_1 = (-1, 0, \dots, 0) \\ \vdots & \vdots \\ e_n = (0, 0, \dots, 1) & \bar{e}_n = (0, 0, \dots, -1) \end{array}$$



$V_2 =$ Del Pezzo surface of degree 6
(over k^{sep})

1. Motivation

2. Derived Categories

3. Arithmetic Toric Varieties

4. Results

Theorem. ([Castravet, Tevelev 2017](#)) $D^b(V_{2n})$ admits a full exceptional collection that is stable under the action of $\text{Gal}(k^{sep}/k)$.

Theorem. ([Klyachko, Voskresenkii 1984](#)) An arithmetic toric variety has a rational point whenever the T -torsor has a rational point. Further, a T -torsor has a rational point if and only if it is the trivial T -torsor.



to find rational points on a toric variety, we just need to detect when the torsor is trivial.

Theorem. (*Ballard, L.*) If X is a generalized Del Pezzo variety, then X has a rational point if and only if $D^b(X)$ admits a full étale exceptional collection.

For forms of dP6, [M. Blunk](#) showed this in 2008!

Example:

$$\text{End}(\mathcal{O}) = \mathbb{C}$$

$$\text{End}(\mathcal{O}(1)) = \mathbb{C}$$



$$D^b(\mathbb{P}_{\mathbb{C}}^1) = \langle \mathcal{O}, \mathcal{O}(1) \rangle$$


$$D^b(C) = \langle \mathcal{O}, \mathcal{F} \rangle$$



$$\mathcal{F} \otimes_{\mathbb{R}} \mathbb{C} \simeq \mathcal{O}(1)^{\oplus 2}$$

$$\text{End}(\mathcal{F}) = \mathbb{H}$$

$\mathbb{H}$ is not isomorphic to $\mathbb{R}$, nor is $\mathbb{H}$ a separable extension of $\mathbb{R}$




desired
Conclusion:

C is not rational!

Proof sketch:

SET-UP

Base field F , V_{2n} defined over $F^{\text{sep.}}$



* Fixing étale F -algebras K, L of appropriate rank gives a T -toric variety " X " which is an F -form of V_{2n} .

* Brauer classes $B \in \text{Br}(K) \cong \text{Br}(L)$ 'detect' when the T -torsor is trivial: B, Q split $\leftrightarrow T$ -torsor is trivial

* Klyachko & Voskresenskii trivial T -torsor $\Rightarrow \exists F$ -rational point on X

Enter: $D^b(X)$.

* Due to [Ballard, Duncan, & McFaddin](#) (2017)  $\text{Gal}(F^{\text{sep}}/F)$ permutes the E_j .

$$D^b(V_{2n}) = \langle E_1, \dots, E_k \rangle$$


 descends to
 $E_i \in D^b(X)$

* It turns out: $\exists i, j$ such that
 $[\text{End}_X(E_i)] \sim [B]$ AND $[\text{End}_X(E_j)] \sim [Q]$

* An exc. collection for $D^b(X)$ yields a decomposition:

$$\mathcal{U}(X) \simeq \bigoplus_{i=1}^k \mathcal{U}(D_i)$$


with $D_i = \text{End}_X(E_i)$!

 Noncommutative motives as a universal additive invariant

No!

Question: Is a full étale exceptional collection a strong enough condition to guarantee the existence of a rational point?

Theorem. ([*Ballard, Duncan, L., McFaddin*](#)) There exists a smooth and geometrically irreducible threefold over $\mathbb{Q}$ with no $\mathbb{Q}$ -points but whose derived category admits a full étale exceptional collection.



Thank
you!